

Macro- and micromorphometric studies of the vascular structures from the Göttingen® minipig

B. Hieb², C. Müller¹, F. Jung², H. Hünigen¹, B. Hamm³, J. Plendl¹, S. M. Niehues³

¹Freie Universität Berlin, Department of Veterinary Medicine, Institute of Veterinary Anatomy, Berlin, Germany;

²Center for Biomaterial Development and Berlin-Brandenburg Center for Regenerative Therapies, Institute of Polymer Research, GKSS Research Center, Teltow, Germany; ³Charité - University Medicine Berlin, Department of Radiology, Berlin, Germany

Applied Cardiopulmonary Pathophysiology 13: 318-321

Key words: minipigs, vasculature, vessel wall

Abstract

Porcine models have become increasingly popular in cardiovascular research, because physiological and anatomical features of the cardiovascular system from pigs including the coronary vascular system and the coronary collateral vessels are comparable to humans. The standard farm pig rapidly increases in body weight and size, potentially confounding serial measurements of cardiac function and morphology. In contrast Göttingen® minipigs have a characteristic growth curve that avoids the dramatic increase in weight in adulthood seen in farm pigs. The Göttingen® minipig is especially suitable for long-term studies because of its inherent small size and ease of handling, even at full maturity, which is reached at 2 y of age compared with 3 y for domestic pigs. However, there is still a need on detailed information about the macro- and microvascular characteristics of the vascular system from the Göttingen® minipigs.

The study was aimed to describe the macro- and microvascular characteristics of adult Göttingen® minipigs (n=18) by use of CT-imaging and histology over a time period of 4 months starting from 16 months of age up to 20 months. The animals showed no clinical symptoms of disease and were kept in-house at a light/dark rhythm of 12:12 under defined climatic conditions. The experiments were licensed by the regional authorities for health and social affairs (LaGeSo), Berlin. The study included the measurement of the length and of the luminal diameter of arteries and veins from the neck, thorax, abdomen, and limbs which are frequently used in experiments with pigs. In addition microscopical and histological parameters (luminal vessel diameter, thickness of the tunica externa, tunica media, and tunica interna) were studied on hematoxylin-eosin-stained sections of the blood vessel.

Introduction

Porcine models have become increasingly popular in cardiovascular research, because physiological and anatomical features of the cardiovascular system of pigs (including the coronary vascular system and the coronary collateral vessels) are quite comparable to humans [1-3]. The standard farm pig rapidly increases in body weight and size, potentially confounding serial measurements of cardiac function and morphology. In contrast, Göttingen® minipigs have a unique growth

characteristic that avoids the dramatic weight and size increase seen in farm pigs even when they reach the adulthood [4, 5]. The Göttingen® minipig was developed in the early 1960s at the Institute of Animal Breeding and Genetics (University of Göttingen, Germany) to reduce space requirements and housing costs for preclinical porcine studies [6]. They were created by crossbreeding the Minnesota minipig with Vietnamese potbelly and German Landrace pigs [7]. The Göttingen® minipig is especially suitable for long-term studies because of its easy handling and inherent small

size even at its full maturity, which is reached in 2 years while it takes 3 years for domestic pigs [8]. However, more detailed information about the macro- and microvascular characteristics of the Göttingen® minipigs has to be elucidated to prove its scientific usefulness.

Materials and methods

Animals, housing and care

In-vivo experiments approved by the regional office for health and social welfare (LaGeSo) of Berlin were performed at the Charité University Clinic, Campus Virchow (Berlin, Germany), Department of Experimental Medicine (certified by ISO 9001) using adult female pigs (n=18) from the Göttingen® Minipig (Ellegaard, Denmark). At the beginning of the test the animals were 17 months old and had a body weight of 26.3 ± 1.8 kg. The animals were cared according to the guidelines of the European societies of laboratory animal sciences. Animals were housed as a group of 6, in an environmentally controlled room (12/12 light/dark-rhythm, 15-24 °C, 55±10 % relative humidity).

Anaesthesia

Prior to CT-imaging and tissue sampling anaesthesia was performed. One night before anaesthesia the animals were fastened, but had free access to water. The minipigs were premedicated by intramuscular (i.m.) injection of 0.5 ml atropine (Atropinum sulfuricum, 1mg/ml, Eifelfango, Germany) and anesthetized by i.m. injection of ketamine (i.m., 27 mg/kg, Ursotamin™, 100 mg/ml, Serumwerk Bernberg, Germany), xylazine (i.m., 3.5 mg/kg, Rompun™ TS, 20 mg/ml, Bayer Vital, Germany) and 3 ml azaperone (i.m., Stresnil™, 40 mg/ml, Janssen Animal Health, Germany). Additionally an electrolyte solution (Iono-steril™, Fresenius, Germany, 1000 ml) was continuously infused intravenously (i.v.).

Computed tomography

The vascular structures of the pigs were studied by computed tomography (CT) analysis at an age of 17, 19 and 21 months according to figure 1. The CT scan was done in prone position on a 64-slice scanner

(LightSpeed 64®; GE Medical Systems, Milwaukee, IL, USA). The scan protocol used contrast-enhancement with automatic intravenous injection of 80 ml nonionic iodinated contrast medium (XenetiX® 350, Guerbet GmbH, Germany 350 mg/ml iodine) in each pig. The scan parameters were standardized (voltage 120 kV, maximal 500 mA with automatic mA-optimization at a noise index of 15, mean mA 490; collimated slice thickness: 64×0.625 mm; total detector width: 55 mm; rotation speed: 0.4 s; table feed per rotation: 55 mm) so the scan speed was approximately 3 s for 30 cm scan length in the z-axis. For volumetric assessment 1.25 mm images were reconstructed without overlapping. Basic image analysis was performed using Advantage Windows 4.2 (GE Medical Systems, Milwaukee, IL, USA) and AccuLite (AccuImage Diagnostics Corporation, South San Francisco, CA, USA).

Histology

Histological examination was done on 5 representative arteries (elastic arteries: Aorta thoracica, Aorta abdominalis, Arteria carotis communis; muscular arteries: A. femoralis, A. renalis) and 5 representative veins (Vena cava cranialis, V. cava caudalis, V. jugularis externa, V. renalis, V. femoralis). After embedding into paraffin, cross sections of these blood vessels with a thickness of 5 µm were made and stained using hematoxylin-eosin. Five slides per sample were prepared and each slide was evaluated at five different fields of view. The thickness of the tunica media and the thickness of the lamina endothelialis were determined using a transmitted light microscope (AxioSkope 2, Zeiss, Germany) and the imaging software AxioVision (Zeiss, Germany).

Statistics

Data are reported as mean value ± standard deviation for continuous variables or as selective frequency for categorical variables and analyzed by Student's t-test or the Chi square test. A p value of less than 0.05 was considered significant.

Results

Body weight

The body weight increased from 26.3 ± 1.8 kg to 33.8 ± 4.7 kg (+ 28.5 %, during the time period between 17 and 21 months of age).

CT analysis

Ct analysis revealed that the luminal diameter and the length of the arteries and veins did not change during the time period between 17 and 21 months of age ($p < 0.05$, figures 2).




Figure 1. Body weight of Göttingen® Minipigs in the age range of 17 to 21 months; (mean value \pm standard deviation, $n = 18$)

Histology

Figure 3 shows the thickness of the tunica media and of the lamina endothelialis from arteries and veins of the animals at an age of 21 months. The thickness of the lamina endothelialis showed no significant difference between the veins and arteries.

Discussion

In order to investigate the properties of implantable biomaterials a biological model is essential before establishing a clinical trial [9]. Therefore, the selection of the appropriate animal model is a major part of experimental work.

The study showed that the vasculature – luminal diameter and the length of the arteries and veins analyzed – remained constant over time in young adult Göttingen® Minipigs despite of a significant increase of the body weight ($p < 0.05$) over 4 months (from month 17 to 21). This is in good agreement with recently published data showing that the diameters of the iliac bifurcation arteries of minipigs (Yucatan) did not change with increasing body weight [10].

In summary the results showed that the adult Göttingen® Minipig proved to be a suitable animal model for in-vivo studies which require long-term stable vascular macro- and micromorphometric parameters, such as implantation of stents, vein cuffs, heart or venous valves and vascular prostheses. The histomorphometric parameters (thickness of the tunica media




Figure 2. Luminal diameter and length of veins and arteries of Göttingen® Minipigs (during the time period between 17 and 21 months of age); CT-based data; (mean value \pm standard deviation, $n = 18$)




Figure 3. Thickness of the tunica media and the lamina endothelialis of arteries and veins of 21 month-old Göttingen® Minipig; data based on histological examination; (mean value \pm standard deviation, $n = 3$)

and the lamina endothelialis) can provide helpful baseline data for developing implants.

References

- Hughes GC, Simons M, Annex BH. Translational physiology: porcine models of human coronary artery disease: implications for preclinical trials of therapeutic angiogenesis. *J Appl Physiol* 2003; 94: 1689-1701
- Grossherr M, Hager A, Gerriets T, Kraatz EG, Misfeld M, Klotz KF. Gelatine and hydroxyl ethyl starch hypervolemic hemodilution – Effect on hemorheology and retinal circulation in a pig model. *Applied Cardiopulmonary Pathophysiology* 2008; 12: 40-46
- Schade I, Knaut M, Plöztze K, Matschke K. Development of an animal model: A key role in investigational research. *Clin Hemorheol Microcirc* 2008; 39: 359-362
- Köhn F, Sharifi AR, Malovrh S, Simianer H. Estimation of genetic parameters for body weight of the Goettingen minipig with random regression models. *J Anim Sci*. 2007; 85 (10): 2423-8
- Köhn F, Sharifi AR, Simianer H. Modeling the growth of the Goettingen minipig. *J Anim Sci* 2007; 85 (1): 84-92
- Beglinger R, Becker M, Eggenberger E, Lombard C. [The Goettingen miniature swine as an experimental animal. 1. Review of literature, breeding and handling, cardiovascular parameters] *Res Exp Med (Berl)*. 1975; 165 (3): 251-63.
- Hashemi SM, Ghods S, Kolodgie FD, Parcham-Azad K, Keane M, Hamamdziec D, Young R, Rippey MK, Virmani R, Litt H, Wilensky RL. A placebo controlled, dose-ranging, safety study of allogenic mesenchymal stem cells injected by endomyocardial delivery after an acute myocardial infarction. *Eur Heart J* 2008; 29 (2): 251-9
- Bollen P, Ellegaard L. The Göttingen minipig in pharmacology and toxicology. *Pharmacol Toxicol*. 1997; 80 (Suppl 2): 3-4
- Rickert D, Scheithauer MO, Coskun S, Kelch S, Lendlein A, Franke RP. The influence of a multifunctional polymeric biomaterial on the concentration of acute phase proteins in an animal model. *Clin Hemorheol Microcirc* 2007; 36: 301-311
- Bushi D, Assaf Y, Grad Y, Nishri B, Yodfat O, Tanne D. Similarity of the swine vasculature to the human carotid bifurcation: analysis of arterial diameters. *J Vasc Interv Radiol*. 2008; 19 (2 Pt 1): 245-51

Address for corresponding: Bernhard Hiebl, M.D., Center for Biomaterial Development, Institute of Polymer Research, GKSS, Research Center, Kantstr. 55, D-14513 Teltow, Germany, bernhard.hiebl@gkss.de